

Phospho-NDRG1-T346 Rabbit pAb

货号: **AYP20382**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	33kDa/35kDa/42kDa
实测分子量	43KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6
细胞定位	Cell membrane,Cytoplasm,Nucleus,centrosome,cytoskeleton,cytosol,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around T346 of human NDRG1 (NP_001128714.1).
序列	SHTSE

靶点信息

研究背景	This gene is a member of the N-myc downregulated gene family which belongs to the alpha/beta hydrolase superfamily. The protein encoded by this gene is a cytoplasmic protein involved in stress responses, hormone responses, cell growth, and differentiation. The encoded protein is necessary for p53-mediated caspase activation and apoptosis. Mutations in this gene are a cause of Charcot-Marie-Tooth disease type 4D, and expression of this gene may be a prognostic indicator for several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
基因ID	10397

基因名	NDRG1
Swiss	Q92597
别名	NDRG1;CAP43;CMT4D;DRG-1;DRG1;GC4;HMSNL;NDR1;NMSL;PROXY1;RIT42;RTP;TARG1;TDD5

产品验证



Western blot analysis of Phospho-NDRG1-T346 expressed in C6 using Phospho-NDRG1-T346 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn