

Phospho-PPIG-S376 Rabbit pAb

货号: **AYP20415**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	40kDa/88kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,C2C12,PC-12
细胞定位	Nucleus matrix,Nucleus speckle
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around S376 of human PPIG (NP_004783.2).
序列	VSSGE

靶点信息

研究背景	PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed:20676357). May be implicated in the folding, transport, and assembly of proteins. May play an important role in the regulation of pre-mRNA splicing.
基因ID	9360
基因名	PPIG

Swiss	Q13427
别名	PPIG;CARS-Cyp;CYP;SCAF10;SRCyp

产品验证



Western blot analysis of Phospho-PPIG-S376 expressed in 293T,C2C12,PC-12 using Phospho-PPIG-S376 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn