

SRA1 Rabbit pAb

货号: **AYP20441**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	25kDa
实测分子量	26kDa, 37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,Mouse testis,Rat liver
细胞定位	cytoplasm,cytosol,intercellular bridge,microtubule cytoskeleton,nucleoplasm,nucleus,plasma membrane, SCAR complex
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human SRA1 (NP_00103031 2.2).
序列	MTRCPAGQAEVEMAELYVKPGNKERGWNDPPQFSYGLQTQAGGPRRSLLTRVAAPQDGSPRPASETSPGPPMGPP PPSSKAPRSPVVGSGPASGVEP

靶点信息

研究背景	Both long non-coding and protein-coding RNAs are transcribed from this gene, and they represent alternatively spliced transcript variants. This gene was initially defined as a non-coding RNA, which is a coactivator for several nuclear receptors (NRs) and is associated with breast cancer. It has now been found that this gene is involved in the regulation of many NR and non-NR activities, including metabolism, adipogenesis and chromatin organization. The long non-coding RNA transcripts interact with a variety of proteins, including the protein encoded by this gene. The encoded protein acts as a transcriptional repressor by binding to the non-coding RNA.
基因ID	10011
基因名	SRA1
Swiss	Q9HD15
别名	SRA1;SRA;SRAP;STRAA1;pp7684

产品验证



Western blot analysis of SRA1 expressed in A-431, Mouse testis, Rat liver using SRA1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn