

— ABLYBIO, Help Your Research



MYCL Rabbit pAb

货号: **AYP20448**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	24kDa/40kDa/43kDa
实测分子量	40KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,BxPC-3
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 40-180 of human MYCL (NP_005367.2).
------	---

靶点信息

研究背景	Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Predicted to be involved in regulation of transcription by RNA polymerase II. Predicted to act upstream of or within regulation of inner ear auditory receptor cell differentiation. Located in chromosome and nucleoplasm.
基因ID	4610
基因名	MYCL
Swiss	P12524 (https://www.uniprot.org/uniprotkb/P12524/entry)
别名	L-Myc,LMYC,MYCL1,bHLHe38,MYCL,MYCL Rabbit pAb,Class E basic helix-loop-helix protein 38,Protein L-Myc-1,V-myc myelocytomatosis viral oncogene homolog

产品验证



Western blot analysis of MYCL expressed in A-549, BxPC-3 using MYCL Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)