

— ABLYBIO, Help Your Research



# WBP2 Rabbit pAb

货号: **AYP20449**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	28kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,A-375,C6,Rat spleen
细胞定位	cytoplasm,cytosol,nucleoplasm,nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 15-140 of human WBP2 (NP_036610.2).
------	---

## 靶点信息

研究背景	The globular WW domain is composed of 38 to 40 semiconserved amino acids shared by proteins of diverse functions including structural, regulatory, and signaling proteins. The domain is involved in mediating protein-protein interactions through the binding of polyproline ligands. This gene encodes a WW domain binding protein that is a transcriptional coactivator of estrogen receptor alpha and progesterone receptor. Defects in this gene have been associated with hearing impairment.
基因ID	23558
基因名	WBP2
Swiss	Q969T9 ( <a href="https://www.uniprot.org/uniprotkb/Q969T9/entry">https://www.uniprot.org/uniprotkb/Q969T9/entry</a> )
别名	WBP2,GRAMD6,WBP-2,WBP2 Rabbit pAb

## 产品验证



Western blot analysis of WBP2 expressed in Rat spleen,U-87MG,A-375 using WBP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)