

— ABLYBIO, Help Your Research



# RRN3 Rabbit pAb

货号: **AYP20459**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	34kDa/53kDa/70kDa/74kDa
实测分子量	34kDa/53kDa/70kDa/74kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,mouse liver,mouse brain
细胞定位	Nucleus,nucleolus
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 550-651 of human RRN3 (NP_060897.3).
------	---

## 靶点信息

研究背景	Required for efficient transcription initiation by RNA polymerase I. Required for the formation of the competent preinitiation complex (PIC. Dissociates from pol I as a consequence of transcription. In vitro, cannot activate transcription in a subsequent transcription reaction (By similarity.
基因ID	54700
基因名	RRN3
Swiss	Q9NYV6 ( <a href="https://www.uniprot.org/uniprotkb/Q9NYV6/entry">https://www.uniprot.org/uniprotkb/Q9NYV6/entry</a> )
别名	RRN3,A-270G1.2,TIFIA,RRN3 homolog,RRN3 Rabbit pAb,Transcription initiation factor IA

## 产品验证



Western blot analysis of RRN3 expressed in HeLa, mouse liver, mouse brain using RRN3 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))