

# NLRC5 Rabbit pAb

货号: **AYP20462**

## 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	204kDa
实测分子量	202kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	mouse lung
细胞定位	centrosome, cytoplasm, cytosol, nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human NLRC5 (NP_115582.4).
序列	MDPVGLQLGNKNLWSCLVRLTKDPEWLNAMKFFLPNTDLDSRNETLDPEQRVILQLNKLHVQGSDTWQSFHCVCM QLEVPLDLEVLLLSTFGYDDGFTSQLGAEKKSQPESQLHHGLKRPHQSCGSSPRRKQCKKQLELAKKYLQLLRTSAQQ RYRSQIPGSGQPHAFHQVYVPPILRRATASLDTPEGAIMGDVKVEDGADVSDLFNTRVKNKGPRVTVLLGKAGMGKTTL AHR

## 靶点信息

研究背景	This gene encodes a member of the caspase recruitment domain-containing NLR family. This gene plays a role in cytokine response and antiviral immunity through its inhibition of NF-kappa-B activation and negative regulation of type I interferon signaling pathways.
基因ID	84166

基因名	NLRC5
Swiss	Q86WI3
别名	NLRC5;CLR16.1;NOD27;NOD4

## 产品验证



Western blot analysis of NLRC5 expressed in mouse lung using NLRC5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)