

# DEPDC6 Rabbit pAb

货号: AYP20476

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100
理论分子量	34kDa/46kDa
实测分子量	34kDa/46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa cells, A2780 cells, COLO 320 cells, HEK-293 cells, MCF-7 cells
细胞定位	
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human DEPDC6 (NP_073620.2).
序列	MEEGGSTGSAGSDSSTSGSGGAQQRELERMAEVLVTGEQLRLRLHEEKVIKDRRHHLKTPNCFVAKELIDWLIHKEAS DRETAIKLMQKLADRGIIHHVCDEHKEFKDVKLFYRFRKDDGTFPLDNEVKAFMRGQRLYEKLMSPENTLLQPREEGVKY ERTFMASEFLDWLVQEGEATTRKEAEQLCHRLMEHGIIQHVS NKHPFVDSNLLYQFRMN

## 靶点信息

研究背景	Negative regulator of the mTORC1 and mTORC2 signaling pathways. Inhibits the kinase activity of both c complexes.
基因ID	64798

基因名	DEPTOR
Swiss	Q8TB45
别名	DEPTOR;DEP.6;DEPDC6

## 产品验证



Western blot analysis of DEPDC6 expressed in HeLa cells,A2780 cells,COLO 320 cells using DEPDC6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 3 0ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time : 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)