

SUPT16H (SPT16) Rabbit pAb

货号: **AYP20484**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	119kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HCT116,Jurkat,C6,Mouse brain
细胞定位	nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 520-695 of human SUPT 16H (SPT16) (NP_009123.1).
序列	KEPHIREMKIYIDKKYETVIMPVFGIATPFHIATIKNISMSVEGDYTYLRINFYCPGSALGRNEGNIFPNPEATFVKEITYRASNIK APGEQTVPALNLQNAFRIIKEVQKRYKTREAEKEKEGIVKQDSLVINLNRSNPKLKDLYIRPNIAQKRMQGSLEAHVNGFRF TSVRGD

靶点信息

研究背景	Transcription of protein-coding genes can be reconstituted on naked DNA with only the general transcript ion factors and RNA polymerase II. However, this minimal system cannot transcribe DNA packaged into c hromatin, indicating that accessory factors may facilitate access to DNA. One such factor, FACT (facilitate s chromatin transcription), interacts specifically with histones H2A/H2B to effect nucleosome disassembly and transcription elongation. FACT is composed of an 80 kDa subunit and a 140 kDa subunit; this gene en codes the 140 kDa subunit.
------	---

基因ID	11198
基因名	SUPT16H
Swiss	Q9Y5B9
别名	CDC68; SPT16; FACTP140; SPT16/CDC68

产品验证



Western blot analysis of SUPT16H (SPT16) expressed in Mouse brain, HeLa, HCT116 using SUPT16H (SPT16) Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn