

# CD52 Rabbit pAb

货号: **AYP20522**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse bone marrow
细胞定位	extracellular region,plasma membrane
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-61 of human CD52 (NP_001794.2).
序列	MKRFLFLLLTISLLVMVQIQTGLSGQNDTSQTSSPSASSNISGGIFLFFVANAIHLFCFS

## 靶点信息

研究背景	CD52 / CDW52 is a small glycosylphosphatidylinositol (GPI) anchored glycoprotein. It has a mature peptide comprising only 12 amino acids and is abundantly expressed on human lymphocytes. From the clinical point of view this protein is an important target for therapeutic interventions aimed at leukocyte depletion in hematological malignancies and post-transplant immunosuppression. CD52 / CDW52 may play a role in carrying and orienting carbohydrate. It is an unusually good target for complement-mediated cell lysis.
基因ID	1043

基因名	CD52
Swiss	P31358
别名	CD52;CDW52;EDDM5

## 产品验证



Western blot analysis of CD52 expressed in Mouse bone marrow using CD52 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)