

# TAOK2 Rabbit pAb

货号: **AYP20638**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	118kDa/119kDa/126kDa/138kDa
实测分子量	138kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	MCF7, 293T
细胞定位	actin cytoskeleton, axon, axonal growth cone, cytoplasm, cytoplasmic vesicle, cytosol, dendritic growth cone, neuron projection, nucleolus, nucleoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 610-730 of human TAO K2 (NP_057235.2).
序列	AEWLLRQKEQLQCCQAEEEEAGLLRRQRQYFELQCRQYKRKMLLARHSLDQDLLREDLNKKQTQKDLECALLLRQHEATR EELRQLQAVQRTRAE LTRLQHQT E LGNQLEYNKRREQLRQ

## 靶点信息

研究背景	This gene encodes a serine/threonine protein kinase that is involved in many different processes, including, cell signaling, microtubule organization and stability, and apoptosis. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
基因ID	9344

基因名	TAOK2
Swiss	Q9UL54
别名	TAOK2;MAP3K17;PSK;PSK1;PSK1-BETA;TAO1;TAO2

## 产品验证



Western blot analysis of TAOK2 expressed in MCF7,293T using TAOK2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)