

HSD17B12 Rabbit pAb

货号: **AYP20701**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	32kD
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	endoplasmic reticulum
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 43-312 of human HSD17B12 (NP_057226.1).
序列	AGVGPGLGEWAVVTGSTDGIGKSYAEELAKHGKVVLSRSKDKLDQVSSEIKEKFKVETRTIAVDFASEDIYDKIKTGLA GLEIGILVNNVGMSSYEYPEYFLDVPDLDNVIKKMININILSVCKMTQLVLPGMVERSKGAILNISSGSGMLPVPLLLTIYSATKTF VDFFSQCLHEEYRSKGVFVQSVLPYFVATKLAKIRKPTLDKPSPEFVKSIAIKTVGLQSRNGLYLIHALMGSIISNLPSWIYLKI VMNMNKSTRAHYLKKTKKN

靶点信息

研究背景	This gene encodes a very important 17beta-hydroxysteroid dehydrogenase (17beta-HSD) that converts estrone into estradiol in ovarian tissue. This enzyme is also involved in fatty acid elongation.
基因ID	51144

基因名	HSD17B12
Swiss	Q53GQ0
别名	HSD17B12;KAR;SDR12C1

产品验证



Western blot analysis of HSD17B12 expressed in human breast cancer using HSD17B12 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn