

NME8 Rabbit pAb

货号: **AYP20703**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	67kDa
实测分子量	67kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,SH-SY5Y,Jurkat
细胞定位	axoneme,cytoplasm,cytosol,nuclear speck,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 299-588 of human NME 8 (NP_057700.3).
序列	FMDAFFPDFKMKMSMKLEKTLALLRPNLFHERKDDVLRRIIKDEDFKILEQRQVVLSEKEAQALCKEYENEDYFNKLIENMTS GPSLALVLLRDNGLQYWKQLLGPRTVEEAIEYFPESLCAQFAMDSLVPVNLQYGSLSLETAEREIQHFFPLQSTLGLIKPHAT SEQREQILKIVKEAGFDLTQVKKMFLTPEQIEKIYPKVTGKDFYKDLLEMLSVGSPSMVILTKWNAVAEWRRLMGPTDPEE AKLLSPDSIRAQFGISKLNIVHGASNAYEAKEVVRNRLFEDPEEN

靶点信息

研究背景	This gene encodes a protein with an N-terminal thioredoxin domain and three C-terminal nucleoside diphosphate kinase (NDK) domains, but the NDK domains are thought to be catalytically inactive. The sea urchin ortholog of this gene encodes a component of sperm outer dynein arms, and the protein is implicated in ciliary function. Mutations in this gene are implicated in primary ciliary dyskinesia type 6.
------	---

基因ID	51314
基因名	NME8
Swiss	Q8N427
别名	NME8;CILD6;HEL-S-99;NM23-H8;SPTRX2;TXNDC3;sptrx-2

产品验证



Western blot analysis of NME8 expressed in U-87MG,SH-SY5Y,Jurkat using NME8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn