

# OSTC Rabbit pAb

货号: **AYP20715**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:100 - 1:500
理论分子量	
实测分子量	23KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F,U-87MG,Mouse liver,Rat liver
细胞定位	Endoplasmic reticulum
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50 to the C-terminus of human OSTC (NP_067050.1).
------	--

## 靶点信息

研究背景	Subunit of the oligosaccharyl transferase (OST complex that catalyzes the initial transfer of a defined glycan (Glc(3Man(9GlcNAc(2 in eukaryotes from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity. May be involved in N-glycosylation of APP (amyloid-beta precursor protein. Can modulate gamma-secretase cleavage of APP by enhancing endoproteolysis of PSEN1.
基因ID	58505

基因名	OSTC
Swiss	Q9NRP0
别名	OSTC;DC2

## 产品验证



Western blot analysis of OSTC expressed in Mouse liver,Rat liver,293F using OSTC Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)