

GORAB Rabbit pAb

货号: **AYP20745**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	21kDa/24kDa/27kDa/44kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,U20S,U-251MG,Mouse spleen
细胞定位	cytoplasm,cytosol,Golgi apparatus,nucleolus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human GORAB (NP_001139511.1).
序列	MSWAAVLAVAAARFGHFHWGCRWPGPMAQGWAGFSEEELRRLKQTKDPFEPQRRLLPAKKSRRQQLQREKALVEQSQKLG LQDGSTSLLEQLLSAPKQRVNV

靶点信息

研究背景	This gene encodes a member of the golgin family, a group of coiled-coil proteins localized to the Golgi. The encoded protein may function in the secretory pathway. The encoded protein, which also localizes to the cytoplasm, was identified by interactions with the N-terminal kinase-like protein, and thus it may function in mitosis. Mutations in this gene have been associated with geroderma osteodysplastica. Alternatively spliced transcript variants have been described.
基因ID	92344

基因名	GORAB
Swiss	Q5T7V8
别名	GORAB;GO;NTKLBP1;SCYL1BP1

产品验证



Western blot analysis of GORAB expressed in Mouse spleen, A-431, U2OS using GORAB Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn