

# NCOA3 Rabbit pAb

货号: **AYP20767**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	147kDa/153kDa/154kDa/155kDa
实测分子量	155kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse brain
细胞定位	cytoplasm,cytosol,extracellular exosome,nucleoplasm,nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human NCOA3 (NP_858045.1).
序列	MSGLGENLDPLASDSRKRKLPDTPGQGLTCSGKRRREQESKYIEELAEELISANLSDIDNFNVKPKCAILKETVRQIRQIK EQGKTISNDDDDVQKADV

## 靶点信息

研究背景	The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Several transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein.
------	---

基因ID	8202
基因名	NCOA3
Swiss	Q9Y6Q9
别名	NCOA3;ACTR;AIB-1;AIB1;CAGH16;CTG26;KAT13B;RAC3;SRC-3;SRC3;TNRC14;TNRC16;TRAM-1;bHLHe42;pCIP

## 产品验证



Western blot analysis of NCOA3 expressed in U-87MG, Mouse brain using NCOA3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)