

FCN3 Rabbit pAb

货号: **AYP20782**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	31kDa/32kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,mouse liver
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 13-288 of human FCN3 (NP_775628.1).
序列	LLLGPPACLKTQEHPSCPGPRELEASKVLLPSCPGAPGSPGKAGPGPQPPGPPGKMGPKGEPGPRNCRELLSQGATL SGWYHLCLPEGRALPVFCMDMTEGGGWLVFQRRQDGSVDFFRSWSSYRAGFGNQEFWLGNEENLHQLTLQGNWEL RVELEDFNGNRTFAHYATFRLGVDHYQLALGKFSEGTAGDSLHSGRPFTTYDADHDSSNSNCAVIVHGAWWYASC YRSNLNGRYAVSEAAAHKYGIDWASGRGVGHPYRRVRMMLR

靶点信息

研究背景	Ficolins are a group of proteins which consist of a collagen-like domain and a fibrinogen-like domain. In human serum, there are two types of ficolins, both of which have lectin activity. The protein encoded by this gene is a thermolabile beta-2-macroglycoprotein found in all human serum and is a member of the ficolin/opsonin p35 lectin family. The protein, which was initially identified based on its reactivity with sera from patients with systemic lupus erythematosus, has been shown to have a calcium-independent lectin activity. The protein can activate the complement pathway in association with MASPs and sMAP, thereby aiding in host defense through the activation of the lectin pathway. Alternative splicing occurs at this locus and two variants, each encoding a distinct isoform, have been identified.
基因ID	8547
基因名	FCN3
Swiss	O75636
别名	FCN3;FCNH;HAKA1;ficolin-3

产品验证



Western blot analysis of FCN3 expressed in A-549, mouse liver using FCN3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn