

SARS-CoV-2 Spike S2 ECD Rabbit pAb

货号: AYP20818

产品信息

反应	SARS-CoV-2
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	DB WB IF/ICC IP
推荐浓度	DB: 1:500 - 1:2000 WB: 1:2000 - 1:6000 IF/ICC: 1:50 - 1:200 IP: 1:2000 - 1:6000
理论分子量	
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Virion membrane
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 800-900 of SARS-CoV-2 Spike S2 ECD (YP_009724390.1).
------	---

靶点信息

研究背景	The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.
基因ID	43740568

基因名	S
Swiss	PODTC2
别名	

产品验证



Western blot analysis of SARS-CoV-2 Spike S2 ECD expressed in 293T using SARS-CoV-2 Spike S2 ECD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/prot eins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Expo sure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn