

FGF17 Rabbit pAb

货号: **AYP20825**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	23kDa/24kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,mouse testis,mouse brain,rat testis,rat brain
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 23-180 of human FGF17 (NP_003858.1).
序列	TQGENHPSPNFNQYVRDQGAMTDQLSRRQIREYQLYSRTSGKHVQVTGRRISATAEDGNKFAKLIVETDTFGSRVRIKGA ESEKYICMNRGKLGKPSGKSKDCVFTEIVLENNYAFQNRHEGWFMFAFTRQGRPRQASRSRQNQREAHFIKRLYQ

靶点信息

研究背景	This gene encodes a member of the fibroblast growth factor (FGF) family. Member of the FGF family possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes including embryonic development cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is expressed during embryogenesis and in the adult cerebellum and cortex and may be essential for vascular growth and normal brain development. Mutations in this gene are the cause of hypogonadotropic hypogonadism 20 with or without anosmia. Alternate splicing results in multiple transcript variants.
------	--

基因ID	8822
基因名	FGF17
Swiss	O60258
别名	FGF17;FGF-13;FGF-17;HH20

产品验证



Western blot analysis of FGF17 expressed in mouse testis, rat testis, SH-SY5Y using FGF17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn