

— ABLYBIO, Help Your Research



# DRG2 Rabbit pAb

货号: **AYP20858**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	40kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat heart,Rat thymus
细胞定位	cytoplasm,cytosol,nucleoplasm,nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 150-260 of human DRG 2 (NP_001379.1).
------	---

## 靶点信息

研究背景	This gene encodes a GTP-binding protein known to function in the regulation of cell growth and differentiation. Read-through transcripts containing this gene and a downstream gene have been identified, but they are not thought to encode a fusion protein. This gene is located within the Smith-Magenis syndrome region on chromosome 17. [provided by RefSeq, Jan 2012]
基因ID	1819
基因名	DRG2
Swiss	P55039 ( <a href="https://www.uniprot.org/uniprotkb/P55039/entry">https://www.uniprot.org/uniprotkb/P55039/entry</a> )
别名	DRG2,DRG2 Rabbit pAb,Translation factor GTPase DRG2

## 产品验证



Western blot analysis of DRG2 expressed in Rat heart,Rat thymus using DRG2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)