

ICA1 Rabbit pAb

货号: **AYP20872**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	
实测分子量	69kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG, BxPC-3
细胞定位	cytoplasm, cytosol, Golgi apparatus, synaptic vesicle membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human ICA1 (NP_001263407).
序列	MSGHKCYPWDLQDRYAQDKSVVNKMQQKYWETKQAFIKATGKKEDEHVVASDADLDAKLELFHSIQRTCLDLSKAIPLY QKRICFLSQEENELGKFLRSQGFQDKTRAGKMMQATGKALCFSSQQLALRNPLCRFHQEVETFRHRAISDTWLTVNRM EQ

靶点信息

研究背景	This gene encodes a protein with an arfaptin homology domain that is found both in the cytosol and as membrane-bound form on the Golgi complex and immature secretory granules. This protein is believed to be an autoantigen in insulin-dependent diabetes mellitus and primary Sjogren's syndrome. Several transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Feb 2013]
------	--

基因ID	3382
基因名	ICA1
Swiss	Q05084
别名	ICA69;ICAp69;ICA1

产品验证



Western blot analysis of ICA1 expressed in U-87MG, BxPC-3 using ICA1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn