

— ABLYBIO, Help Your Research



EXOSC10 Rabbit pAb

货号: **AYP20887**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	98kDa/100kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Raji,Mouse testis,Mouse spleen,Mouse lung,Rat brain
细胞定位	cytoplasm,cytoplasmic exosome (RNase complex),cytosol,nuclear exosome (RNase complex),nucleolar exosome (RNase complex),nucleolus,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 586-885 of human EXOSC10 (NP_001001998.1).
------	--

靶点信息

研究背景	Putative catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. EXOSC10 has 3'-5' exonuclease activity (By similarity. EXOSC10 is required for nucleolar localization of C1D and probably mediates the association of MTREX, C1D and MPHOSPH6 with the RNA exosome involved in the maturation of 5.8S rRNA.
基因ID	5394
基因名	EXOSC10
Swiss	Q01780 (https://www.uniprot.org/uniprotkb/Q01780/entry)
别名	EXOSC10,PM-Scl,PM/Scl-100,PMSC1,PMSC2,RRP6,Rrp6p,p2,p3,p4,EXOSC10 Rabbit pAb,Autoantigen PM/Scl 2,P100 polymyositis-scleroderma overlap syndrome-associated autoantigen,Polymyositis/scleroderma autoantigen 100 kDa,Polymyositis/scleroderma autoantigen 2

产品验证



Western blot analysis of EXOSC10 expressed in Mouse testis,Rat brain,A-549 using EXOSC10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)