

— ABLYBIO, Help Your Research



# TNFAIP1 Rabbit pAb

货号: **AYP20905**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	24kDa/36kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	cytoplasm, cytosol, endosome, nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 90-160 of human TNFAIP1 (NP_066960.1).
------	--

## 靶点信息

研究背景	This gene was identified as a gene whose expression can be induced by the tumor necrosis factor alpha (TNF) in umbilical vein endothelial cells. Studies of a similar gene in mouse suggest that the expression of this gene is developmentally regulated in a tissue-specific manner. [provided by RefSeq, Jul 2008]
基因ID	7126
基因名	TNFAIP1
Swiss	Q13829 ( <a href="https://www.uniprot.org/uniprotkb/Q13829/entry">https://www.uniprot.org/uniprotkb/Q13829/entry</a> )
别名	B12,B61,BTBD34,EDP1,hBACURD2,TNFAIP1,TNFAIP1 Rabbit pAb,BTB/POZ domain-containing protein TNFAIP1,Protein B12,Tumor necrosis factor,alpha-induced protein 1,endothelial,BACURD2

## 产品验证



Western blot analysis of TNFAIP1 expressed in K-562 using TNFAIP1 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)