

ZNF16 Rabbit pAb

货号: **AYP20909**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	77kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	OVCAR3, SKOV3, A-549, NCI-H460
细胞定位	nucleolus, nucleoplasm, nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-210 of human ZNF16 (NP_008889.2).
序列	MPSLRTRREEAEMELSVPGSPWTPAAQARVRDAPAVTHPGSAACGTPCCSDTELEAICPHYQQPDCDTRTEDKEFLHKE DIHEDLESQAEISENYAGDVSQVPELGDLCDDVSRDWGVPEGRRLPQSLSQEGDFTPAAMGLLRGPLGEKDLDCNGF DSRFSLSPLNMACQEIPTEERPHPYDMGGQSFQHSVDLTGHEGVPTAESPLI

靶点信息

研究背景	The protein encoded by this gene contains multiple tandem zinc finger motifs. The encoded protein is involved in the differentiation of erythroid and megakaryocytic cells. This gene is located in a cluster of related genes on chromosome 8 encoding zinc finger proteins. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2012]
基因ID	7564

基因名	ZNF16
Swiss	P17020
别名	HZF1;KOX9;ZNF16

产品验证



Western blot analysis of ZNF16 expressed in OVCAR3,SKOV3,A-549 using ZNF16 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn