

— ABLYBIO, Help Your Research



RBM7 Rabbit pAb

货号: **AYP20949**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Jurkat,Mouse thymus
细胞定位	nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-266 of human RBM7 (NP_057174.1).
------	------------------------------------------------------------------------------------------------------------------

靶点信息

研究背景	RNA-binding subunit of the trimeric nuclear exosome targeting (NEXT complex, a complex that functions as an RNA exosome cofactor that directs a subset of non-coding short-lived RNAs for exosomal degradation. NEXT is involved in surveillance and turnover of aberrant transcripts and non-coding RNAs. Binds preferentially polyuridine sequences and associates with newly synthesized RNAs, including pre-mRNAs and short-lived exosome substrates such as promoter upstream transcripts (PROMPTs, enhancer RNAs (eRNAs, and 3'-extended products from small nuclear RNAs (snRNAs. Participates in several biological processes including DNA damage response (DDR and stress response. During stress response, activation of the p38 MAPK-MK2 pathway decreases RBM7-RNA-binding and subsequently the RNA exosome degradation activities, thereby modulating the turnover of non-coding transcriptome. Participates in DNA damage response (DDR, through its interaction with MEPCE and LARP7, the core subunits of 7SK snRNP complex, that release the positive transcription elongation factor b (P-TEFb complex from the 7SK snRNP. In turn, activation of P-TEFb complex induces the transcription of P-TEFb-dependent DDR genes to promote cell viability.
基因ID	10179
基因名	RBM7
Swiss	Q9Y580 (https://www.uniprot.org/uniprotkb/Q9Y580/entry)
别名	RBM7,RBM7 Rabbit pAb,RNA-binding motif protein 7

产品验证



Western blot analysis of RBM7 expressed in A-549, Jurkat, Mouse thymus using RBM7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)