

— ABLYBIO, Help Your Research



# SLU7 Rabbit pAb

货号: **AYP20953**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis,Mouse lung,Rat spleen,Rat lung
细胞定位	cytoplasm,cytosol,nuclear speck,nucleoplasm,nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human SLU7 (N P_006416.3).
------	---

## 靶点信息

研究背景	Pre-mRNA splicing occurs in two sequential transesterification steps. The protein encoded by this gene is a splicing factor that has been found to be essential during the second catalytic step in the pre-mRNA splicing process. It associates with the spliceosome and contains a zinc knuckle motif that is found in other splicing factors and is involved in protein-nucleic acid and protein-protein interactions. [provided by RefSeq, Jul 2008]
基因ID	10569
基因名	SLU7
Swiss	O95391 ( <a href="https://www.uniprot.org/uniprotkb/O95391/entry">https://www.uniprot.org/uniprotkb/O95391/entry</a> )
别名	9G8,hSlu7,SLU7,SLU7 Rabbit pAb

## 产品验证



Western blot analysis of SLU7 expressed in Mouse testis,Rat spleen,Mouse lung using SLU7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))