

WBP4 Rabbit pAb

货号: **AYP20972**

产品信息

| | |
|-------|--|
| 反应 | Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | |
| 实测分子量 | 43kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Rat thymus |
| 细胞定位 | nuclear speck, nucleoplasm, nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-376 of human WBP4 (NP_009118.1). |
| 序列 | MADYWKSQPKKFCDYCKCWIADNRPSVEFHERGKNHKENVAKRISEIKQKSLDKAKEEEKASKEFAAMEAAALKAYQEDL KRLGLESEILEPSITPVTSTIPPTSTSNQQKEKKEKKRKKDPSKGRWVEGITSEGYHYYDLISGASQWEKPEGFQGDLKK TAVKTVWVEGLSEDFGFTYYNTETGESRWEKPDFFIPHTSDLPSSKVNENSLGTLDESKSSDSHSDSDGEQAEEGGVS TETEKPKIKFKEKNKNSDGGSDPETQKEKSIQKQNSLGSNEEKSKTLKKSNPYGEWQEIKQEVESHEEVDLELPSTENEYV STSEADGGGEPKVVFEKKTVSLGVMADGVAPVFKRRRTENGKSRNLRQGGDDQ |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes WW domain-containing binding protein 4. The WW domain represents a small and compact globular structure that interacts with proline-rich ligands. This encoded protein is a general spliceosomal protein that may play a role in cross-intron bridging of U1 and U2 snRNPs in the spliceosomal complex A. [provided by RefSeq, Jul 2008] |
|------|--|

| | |
|-------|------------|
| 基因ID | 11193 |
| 基因名 | WBP4 |
| Swiss | O75554 |
| 别名 | FBP21;WBP4 |

产品验证



Western blot analysis of WBP4 expressed in Rat thymus using WBP4 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn