

HSPA4L Rabbit pAb

货号: **AYP20975**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	
实测分子量	115kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,K-562,HeLa,Mouse brain,Rat testis,Rat brain
细胞定位	cytoplasm,cytosol,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 500-600 of human HSPA4L (NP_055093.2).
序列	KQNLEGDHSDAPMETETSFKNENKDNMDKMQVDQEEGHQKCHAHTPEEEIDHTGAKTKSAVSDKQDRNLNQLTKKGK VKSIDLPIQSSLCRQLGQDLLNSY

靶点信息

研究背景	The protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015]
基因ID	22824

基因名	HSPA4L
Swiss	O95757
别名	APG-1;HSPH3;Osp94;HSPA4L

产品验证



Western blot analysis of HSPA4L expressed in Mouse brain,Rat testis,U-87MG using HSPA4L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn