

MBIP Rabbit pAb

货号: **AYP21024**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	32kDa/34kDa/39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	cytosol, mitotic spindle, nucleolus, nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-344 of human MBIP (N P_057670.2).
序列	MAAATELNRPSSGDRNLERRCRPNLSREVLVEIFRSLHTLVGQLDLRDDVVKITIDWNKLQSLSAFQPALLFSALEQHILYLQ PFLAKLQSPIKEENTTAVEEIGRTEMGNKNEVNDKFSIGDLQEEKHKESDLRDVKKTIHFDPPEVVQIKAGKAEIDRRISAFI ERKQAEINENNVRFCNVIDCNQENSCARTDAIFTYPYPGFKSHVKVSRVVNTYGPQTRPEGIPGSGHKPNSMLRDCGNQA VEERLQNI EAHLRLQTGGPVPRDIYQRIKKLEDKILELEGISPEYFQSVSFGKRRKVPQPPQNYSLAELDEKISALKQALLRK SREAESMATHHLP

靶点信息

研究背景	Inhibits the MAP3K12 activity to induce the activation of the JNK/SAPK pathway. Component of the ATAC complex, a complex with histone acetyltransferase activity on histones H3 and H4.
基因ID	51562

基因名	MBIP
Swiss	Q9NS73
别名	MBIP

产品验证



Western blot analysis of MBIP expressed in Rat brain using MBIP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn