

DHX29 Rabbit pAb

货号: **AYP21026**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	155kDa
实测分子量	155kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1020-1369 of human D HX29 (NP_061903.2).
序列	LHIMKCNLGSPEDFLSKALDPPQLQVISNAMNLLRKIGACELNEPKLTPLGQHLAALPVNVKIGKMLIFGAIFGCLDPVATLA AVMTEKSPFTTPIGRKDEADLAKSALAMADSDHLTIYNAYLGWKKARQEGGYRSEITYCRRNFLNRTSLLTLEDVKQELIKL VKAAGFSSSTTSTSWEGNRASQTLFQEIALLKAVLVAGLYDNVVGKIYTKSVDVTEKLACIVETAQGKAQVHPSSVNRDL QTHGWLLYQEKIRYARVYLRETTLITPPVLLFGGDIEVQHRERLLSIDGWYIFQAPVKIAVIFKQLRVLIDSVLRKKLENPKMS LENDKILQIITELIKTENN

靶点信息

研究背景	This gene encodes a member of the DEAH (Asp-Glu-Ala-His) subfamily of proteins, part of the DEAD (Asp-Glu-Ala-Asp) box family of RNA helicases. The encoded protein functions in translation initiation, and is specifically required for ribosomal scanning across stable mRNA secondary structures during initiation codon selection. This protein may also play a role in sensing virally derived cytosolic nucleic acids. Knockdown of this gene results in reduced protein translation and impaired proliferation of cancer cells. [provided by RefSeq, Sep 2016]
基因ID	54505
基因名	DHX29
Swiss	Q7Z478
别名	DDX29;DHX29

产品验证



Western blot analysis of DHX29 expressed in HepG2 using DHX29 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn