

# MTPAP Rabbit pAb

货号: **AYP21032**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,U-251MG,Jurkat,Mouse spleen
细胞定位	mitochondrion,nucleoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 383-582 of human MTPAP (NP_060579.3).
序列	MVIFFLQRRSPILPTLDSLKTLADAEDKCVIEGNNCTFVRDLSRIKPSQNTETLELLLKEFFEYFGNFAFDKNSINIRQGREQ NKPDSSPLYIQNPFETSLNISKNVSQLKQFVDLARESAWILQQEDTDRPSISSNRPWGLVSLLLPSAPNRKSFTKKKSN KFAIETVKNLLESLKGNRTENFTKTSKRTISTQT

## 靶点信息

研究背景	The protein encoded by this gene is a member of the DNA polymerase type-B-like family. This enzyme synthesizes the 3' poly(A) tail of mitochondrial transcripts and plays a role in replication-dependent histone mRNA degradation.[provided by RefSeq, Jan 2011]
基因ID	55149

基因名	MTPAP
Swiss	Q9NVV4
别名	PAPD1;SPAX4;MTPAP

## 产品验证



Western blot analysis of MTPAP expressed in Mouse spleen,U-87MG,U-251MG using MTPAP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)