

# NDC1 Rabbit pAb

货号: **AYP21047**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	76kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Jurkat,HeLa,Mouse heart,Mouse brain
细胞定位	actin cytoskeleton,cytoplasm,nuclear envelope,nuclear membrane,nuclear pore,nuclear pore transmembrane ring,plasma membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 340-530 of human NDC 1 (NP_060557.3).
序列	QYSPSRRQEVFSLSQPGGHPHNWTAISRECLNLLNGMTQKLILYQEEAATNGRVSSSYVPEPKKLSPEETAFTQPKSSQMPRPSVPLVKTSLSFKLSTPDVVSPFGTPFGSSVMNRMAGIFDVNTCYGSPQSPQLIRRGPRWLWTSASDQMQMTEFSNPSPSTSISAEGKTMRQPSVIYSWIQNKRETIK

## 靶点信息

研究背景	Component of the nuclear pore complex (NPC, which plays a key role in de novo assembly and insertion of NPC in the nuclear envelope. Required for NPC and nuclear envelope assembly, possibly by forming a link between the nuclear envelope membrane and soluble nucleoporins, thereby anchoring the NPC in the membrane.
------	--

基因ID	55706
基因名	NDC1
Swiss	Q9BTX1
别名	NET3;TMEM48;NDC1

## 产品验证



Western blot analysis of NDC1 expressed in Mouse heart, MCF7, Jurkat using NDC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)