

DCAF17 Rabbit pAb

货号: **AYP21083**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	59kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Rat spleen, Rat lung
细胞定位	cytosol, nucleolus, nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 235-335 of human DCAF17 (NP_079276.2).
序列	LYSFQTIAEQFMQQKLDLGCACRWGGTTGTVGEAPFGIPCNIKITDMPPLLFEVSSLENAFQIGGHPWHYIVTPNKKKQKG VFHICALKDNSLAKNGIQEM

靶点信息

研究背景	This gene encodes a nuclear transmembrane protein that associates with cullin 4A/damaged DNA binding protein 1 ubiquitin ligase complex. Mutations in this gene are associated with Woodhouse-Sakati syndrome. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009]
基因ID	80067

基因名	DCAF17
Swiss	Q5H9S7
别名	C20orf37;C2orf37;DCAF17

产品验证



Western blot analysis of DCAF17 expressed in Rat spleen, Rat lung using DCAF17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn