

PGLYRP2 Rabbit pAb

货号: **AYP21109**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	62kDa/68kDa
实测分子量	62kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	extracellular exosome,extracellular region
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 300-400 of human PGLYRP2 (NP_443122.3).
序列	LSQYYGAGVARDPGRFSNFRRQNGAALTSASILAQQVWGLVLLQRLEPVHLQLQCMSQEQLAQVAANATKEFTEAFLGCPAIHPRCRWGAAPYRGRPKLL

靶点信息

研究背景	This gene encodes a peptidoglycan recognition protein, which belongs to the N-acetylmuramoyl-L-alanine amidase 2 family. This protein hydrolyzes the link between N-acetylmuramoyl residues and L-amino acid residues in bacterial cell wall glycopeptides, and thus may play a scavenger role by digesting biologically active peptidoglycan into biologically inactive fragments. [provided by RefSeq, Sep 2011]
基因ID	114770

基因名	PGLYRP2
Swiss	Q96PD5
别名	HMFT0141;PGLYRPL;PGRP-L;PGRPL;TAGL-like;tagL;tagL-alpha;tagI-beta;PGLYRP2

产品验证



Western blot analysis of PGLYRP2 expressed in Mouse liver using PGLYRP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn