

# OSBPL8 Rabbit pAb

货号: **AYP21111**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	
实测分子量	101kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen,Mouse kidney,Rat thymus,Rat kidney
细胞定位	cortical endoplasmic reticulum,cytosol,endoplasmic reticulum,nuclear membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-110 of human OSBPL8 (NP_065892.1).
序列	MEGGLADGEPDRTSLLGDSKDVLGPSTVVANSDESQLLTPGKMSQRQGKEAYPTPTKDLHQPSLSPASPHSQGFERGGK EDISQNKDESSLMSKSKSESKLYNGSEKDSS

## 靶点信息

研究背景	This gene encodes a member of a family of proteins containing an N-terminal pleckstrin homology domain and a highly conserved C-terminal oxysterol-binding protein-like sterol-binding domain. It binds multiple lipid-containing molecules, including phosphatidylserine, phosphatidylinositol 4-phosphate (PI4P) and oxysterol, and promotes their exchange between the endoplasmic reticulum and the plasma membrane. An alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016]
------	--

基因ID	114882
基因名	OSBPL8
Swiss	Q9BZF1
别名	MST120;MSTP120;ORP8;OSBP10;OSBPL8

## 产品验证



Western blot analysis of OSBPL8 expressed in Mouse spleen,Rat thymus,Mouse kidney using OSBPL8 R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)