

STK35 Rabbit pAb

货号: AYP21122

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	58kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney
细胞定位	cytoplasm, nuclear body, nucleolus, nucleoplasm, nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 190-380 of human STK35 (NP_543026.2).
序列	RPEGGGGSARPRYSLLAEIGRGSYGVVYEAVAGRSGARVAVKKIRCDAPENVELALAEFWALTSKRRRHQNVVQFEECVLQRNGLAQRMSHGKSSQLYLRVETSLKGERILGYAEPCYLWFMFCEGGDLNQYVLSRRPDPATNKSFMLQLTSAIAFLHKNHIVHRDLKPDNILITERSGTPILKVA

靶点信息

研究背景	The protein encoded by this gene is a kinase that is predominantly found in the nucleus. However, it can interact with PDLIM1/CLP-36 in the cytoplasm and localize to actin stress fibers. The encoded protein may be a regulator of actin stress fibers in nonmuscle cells. [provided by RefSeq, Jul 2008]
基因ID	140901

基因名	STK35
Swiss	Q8TDR2
别名	CLIK1;STK35L1;STK35

产品验证



Western blot analysis of STK35 expressed in Mouse kidney using STK35 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn