

# NAF1 Rabbit pAb

货号: **AYP21125**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	62-70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,Mouse testis
细胞定位	cytoplasm,nucleoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 150-400 of human NAF1 (NP_612395.2).
序列	PPVLSGDGDDDLQIEKENKNFPLKTKDELLLNELPSVEELTIILPEDIELKPLGMVSSIIQLVIIESMTNLPPVNEETVIFKSDRQ AAGKIFEIFGPVAHPFYVLRFNSSDHIESKGIKIKETMYFAPSMKDFTQYIFTEKLKQDKGSDASWKNDQEPPEALDFSD EKEKEAKQRKKSQIQGRKLLKSEFNPEGDFTEVHQNWNNAHSSASEHAKGYRNREFTRGFSRARYPRSCHGRPPPQHFY NSEH

## 靶点信息

研究背景	RNA-binding protein required for the maturation of box H/ACA snoRNPs complex and ribosome biogenesis . During assembly of the H/ACA snoRNPs complex, it associates with the complex and disappears during maturation of the complex and is replaced by NOLA1/GAR1 to yield mature H/ACA snoRNPs complex. Pro bably competes with NOLA1/GAR1 for binding with DKC1/NOLA4.
------	--

基因ID	92345
基因名	NAF1
Swiss	Q96HR8
别名	NAF1

## 产品验证



Western blot analysis of NAF1 expressed in K-562, Mouse testis using NAF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)