

— ABLYBIO, Help Your Research



# CHODL Rabbit pAb

货号: **AYP21127**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis,Rat spleen
细胞定位	centrosome,cytoplasm,cytosol,perinuclear region of cytoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 110-210 of human CHODL (NP_001191103.1).
------	--

## 靶点信息

研究背景	This gene encodes a type I membrane protein with a carbohydrate recognition domain characteristic of C-type lectins in its extracellular portion. In other proteins, this domain is involved in endocytosis of glycoproteins and exogenous sugar-bearing pathogens. This protein localizes predominantly to the perinuclear region. Several transcript variants encoding a few different isoforms have been found for this gene. [provided by RefSeq, Feb 2011]
基因ID	140578
基因名	CHODL
Swiss	Q9H9P2 ( <a href="https://www.uniprot.org/uniprotkb/Q9H9P2/entry">https://www.uniprot.org/uniprotkb/Q9H9P2/entry</a> )
别名	C21orf68,MT75,PRED12,CHODL,CHODL Rabbit pAb,Transmembrane protein MT75

## 产品验证



Western blot analysis of CHODL expressed in Mouse testis,Rat spleen using CHODL Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)