

# ZC3H12D Rabbit pAb

货号: **AYP21149**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	25kDa/36kDa/58kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,THP-1,Mouse spleen,Mouse liver,Rat spleen
细胞定位	cytoplasm,cytoplasmic ribonucleoprotein granule,nucleoplasm,nucleus,P-body
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 150-230 of human ZC3H12D (NP_997243.2).
序列	QHVLAEALERQAVLVYTPSRKVHKGKRLVCYDDRYIVKVAYEQDGVIVSNDNYRDLQSENPEWKWFIEQRLLMFSFVNDRFMP

## 靶点信息

研究背景	May regulate cell growth likely by suppressing RB1 phosphorylation. May function as RNase and regulate the levels of target RNA species (Potential. In association with ZC3H12A enhances the degradation of interleukin IL-6 mRNA level in activated macrophages. Serve as a tumor suppressor in certain leukemia cells . Overexpression inhibits the G1 to S phase progression through suppression of RB1 phosphorylation.
基因ID	340152

基因名	ZC3H12D
Swiss	A2A288
别名	C6orf95;MCPIP4;TFL;dJ281H8.1;p34;ZC3H12D

## 产品验证



Western blot analysis of ZC3H12D expressed in Mouse spleen, Rat spleen, U-87MG using ZC3H12D Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)