

— ABLYBIO, Help Your Research



RAB7B Rabbit pAb

货号: **AYP21150**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | |
| 实测分子量 | 23kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | A-549,Mouse testis,Mouse brain |
| 细胞定位 | Golgi apparatus,late endosome,lysosome,phagocytic vesicle,trans-Golgi network |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-199 of human RAB7B (NP_001157994.1). |
|------|--|

靶点信息

| | |
|-------|--|
| 研究背景 | Controls vesicular trafficking from endosomes to the trans-Golgi network (TGN). Acts as a negative regulator of TLR9 signaling and can suppress TLR9-triggered TNFA, IL6, and IFNB production in macrophages by promoting TLR9 lysosomal degradation. Also negatively regulates TLR4 signaling in macrophages by promoting lysosomal degradation of TLR4. Promotes megakaryocytic differentiation by increasing NF-kappa-B-dependent IL6 production and subsequently enhancing the association of STAT3 with GATA1. Not involved in the regulation of the EGF- and EGFR degradation pathway. |
| 基因ID | 338382 |
| 基因名 | RAB7B |
| Swiss | Q96AH8 (https://www.uniprot.org/uniprotkb/Q96AH8/entry) |
| 别名 | RAB7,RAB7B,RAB7B Rabbit pAb |

产品验证



Western blot analysis of RAB7B expressed in A-549, Mouse testis, Mouse brain using RAB7B Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)