

Ubiquityl-Histone H2A-K119 Rabbit pAb

货号: **AYP21169**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	23KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6
细胞定位	extracellular exosome,nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic ubiquitinated peptide around K119 of human HIST1H2AB (NP_734466.1).
------	---

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures.H2A clustered histone (H2AC) consists of several H2A family members, which function as replication-dependent histone proteins.
基因ID	3012,8335
基因名	H2AC4, H2AC8

Swiss	P04908
别名	HIST1H2AB

产品验证



Western blot analysis of Ubiquityl-Histone H2A-K119 expressed in C6 using Ubiquityl-Histone H2A-K119 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn