

— ABLYBIO, Help Your Research



MED15 Rabbit pAb

货号: **AYP21188**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	87kDa
实测分子量	87kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart,Mouse brain,Rat heart,Rat brain
细胞定位	cytoplasm,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human MED15 (NP_001003 891.1).
------	--

靶点信息

研究背景	The protein encoded by this gene is a subunit of the multiprotein complexes PC2 and ARC/DRIP and may function as a transcriptional coactivator in RNA polymerase II transcription. This gene contains stretches of trinucleotide repeats and is located in the chromosome 22 region which is deleted in DiGeorge syndrome. Alternative splicing results in multiple transcript variants.
基因ID	51586
基因名	MED15
Swiss	Q96RN5 (https://www.uniprot.org/uniprotkb/Q96RN5/entry)
别名	MED15,ARC105,CAG7A,CTG7A,PCQAP,TIG-1,TIG1,TNRC7,MED15 Rabbit pAb,Activator-recruited cofactor 1 05 kDa component,CTG repeat protein 7a,Mediator complex subunit 15,Positive cofactor 2 glutamine/Q-rich-associated protein,TPA-inducible gene 1 protein,Trinucleotide repeat-containing gene 7 protein

产品验证



Western blot analysis of MED15 expressed in Mouse heart,Rat heart,Mouse brain using MED15 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)