

— ABLYBIO, Help Your Research



TDP2 Rabbit pAb

货号: **AYP21211**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	32kDa/40kDa/44kDa
实测分子量	41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Nucleus,PML body,nucleolus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human TDP2 (NP_057698.2).
------	--

靶点信息

研究背景	This gene encodes a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor associated factors (TRAFs), and inhibits nuclear factor-kappa-B activation. This protein has sequence and structural similarities with APE1 endonuclease, which is involved in both DNA repair and the activation of transcription factors.
基因ID	51567
基因名	TDP2
Swiss	O95551 (https://www.uniprot.org/uniprotkb/O95551/entry)
别名	TDP2,AD022,EAP2,EAPII,TTRAP,dj30M3.3,hTDP2,TDP2 Rabbit pAb,5'-tyrosyl-DNA phosphodiesterase,ETS1-associated protein 2,ETS1-associated protein II,TRAF and TNF receptor-associated protein,Tyrosyl-RNA phosphodiesterase,VPg unlinkase

产品验证



Western blot analysis of TDP2 expressed in HeLa using TDP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)