

— ABLYBIO, Help Your Research



FBXW8 Rabbit pAb

货号: **AYP21297**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | |
| 实测分子量 | 67kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-87MG,A-549,A-431,Rat lung,Rat heart |
| 细胞定位 | cytosol,Golgi apparatus,perinuclear region of cytoplasm |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 500-598 of human FBXW8 (NP_699179.2). |
|------|--|

靶点信息

| | |
|-------|---|
| 研究背景 | This gene encodes a member of the F-box protein family, members of which are characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into three classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene contains a WD-40 domain, in addition to an F-box motif, so it belongs to the Fbw class. Alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008] |
| 基因ID | 26259 |
| 基因名 | FBXW8 |
| Swiss | Q8N3Y1 (https://www.uniprot.org/uniprotkb/Q8N3Y1/entry) |
| 别名 | FBW6,FBW8,FBX29,FBXO29,FBXW6,FBXW8,FBXW8 Rabbit pAb,F-box and WD-40 domain-containing protein 8,F-box only protein 29 |

产品验证



Western blot analysis of FBXW8 expressed in Rat lung,U-87MG,A-549 using FBXW8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)