

# HSPB11 Rabbit pAb

货号: **AYP21301**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	16kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,293T,Mouse kidney,Mouse brain,Rat thymus
细胞定位	centrosome,ciliary tip,cilium
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-144 of human HSPB11 (NP_057210.2).
序列	MRKIDLCLSSEGSEVILATSSDEKHPPENIIDGNPETFWTTTGMFPQEFICFHKHVRIERLVIQSYFVQTLKIEKSTSKEPVD FEQWIEKDLVHTEGQLQNEEIVAHDG SATYLRFIIVSAFDHFASVHSVSAEGTVVSNLSS

## 靶点信息

研究背景	Component of the IFT complex B required for sonic hedgehog/SHH signaling. May mediate transport of SHH components: required for the export of SMO and PTCH1 receptors out of the cilium and the accumulation of GLI2 at the ciliary tip in response to activation of the SHH pathway, suggesting it is involved in the dynamic transport of SHH signaling molecules within the cilium. Not required for ciliary assembly. Its role in intraflagellar transport is mainly seen in tissues rich in ciliated cells such as kidney and testis. Essential for male fertility, spermiogenesis and sperm flagella formation. Plays a role in the early development of the kidney. May be involved in the regulation of ureteric bud initiation (By similarity).
基因ID	51668
基因名	HSPB11
Swiss	Q9Y547
别名	HSPB11;C1orf41;HSPCO34;IFT25;PP25

## 产品验证



Western blot analysis of HSPB11 expressed in Mouse kidney, Rat thymus, A-549 using HSPB11 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)