

MRPL55 Rabbit pAb

货号: **AYP21376**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	15kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	mitochondrial inner membrane, mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-128 of human MRPL55 (NP_001308213.1).
序列	MAAVGSLGRLRQSTVKATG PALRRLLHTSSWRADSSRASLTRVHRQAYARLYPVLLVKQD GSTIHIRYREPRRMLAMPIDL DTLSPEERRARLRKREAQLQSRKEYEQELSDDLHVERYRQFWTRTKK

靶点信息

研究背景	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. Multiple transcript variants encoding two different isoforms were identified through sequence analysis.
基因ID	128308
基因名	MRPL55
Swiss	Q7Z7F7
别名	L55nt;MRP-L55;AAVG5835;PRO19675;MRPL55

产品验证



Western blot analysis of MRPL55 expressed in MCF7 using MRPL55 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn