

— ABLYBIO, Help Your Research



# CXCL10/IP-10 Rabbit pAb

货号: **AYP21424**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:1000 - 1:5000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	10kDa
实测分子量	11KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1
细胞定位	Secreted
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 22-98 of human CXCL10/IP-10 (NP_001556.2).
------	--

## 靶点信息

研究背景	This antimicrobial gene encodes a chemokine of the CXC subfamily and ligand for the receptor CXCR3. Binding of this protein to CXCR3 results in pleiotropic effects, including stimulation of monocytes, natural killer and T-cell migration, and modulation of adhesion molecule expression.
基因ID	3627
基因名	CXCL10
Swiss	P02778 ( <a href="https://www.uniprot.org/uniprotkb/P02778/entry">https://www.uniprot.org/uniprotkb/P02778/entry</a> )
别名	CXCL10,C7,IFI10,INP10,IP-10,SCYB10,crg-2,gIP-10,mob-1,CXCL10/IP-10 Rabbit pAb,10 kDa interferon gamma-induced protein,Small-inducible cytokine B10

## 产品验证



Western blot analysis of CXCL10/IP-10 expressed in THP-1 using CXCL10/IP-10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)