

# GCN1L1 Rabbit pAb

货号: **AYP21476**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	292kDa
实测分子量	275KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	cytoplasm,cytosol
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 2462-2671 of human GCN1L1 (NP_006827.1).
序列	LSAVLQQLLADVSGIDWMVVRHGRSLALSVAVNVAPGRLCAGRYSSDVQEMILSSATADRIPIAVSGVVRMGFLMRHHIE TGGGQLPAKLSSLFVKCLQNPSSDIRLVAEKMIWWANKDPLPPLDPQAIKPKLALLDNTKDKNTVVRAYSDAQIVNLLKM RQGEEVFQSLSKILDVASLEVLNEVNRRLKSLKSLASQADSTEQVDDTILT

## 靶点信息

研究背景	Acts as a positive activator of the EIF2AK4/GCN2 protein kinase activity in response to amino acid starvation. Forms a complex with EIF2AK4/GCN2 on translating ribosomes; during this process, GCN1 seems to act as a chaperone to facilitate delivery of uncharged tRNAs that enter the A site of ribosomes to the tRNA-binding domain of EIF2AK4/GCN2, and hence stimulating EIF2AK4/GCN2 kinase activity. Participates in the repression of global protein synthesis and in gene-specific mRNA translation activation, such as the transcriptional activator ATF4, by promoting the EIF2AK4/GCN2-mediated phosphorylation of eukaryotic translation initiation factor 2 (eIF-2-alpha/EIF2S1 on 'Ser-52', and hence allowing ATF4-mediated reprogramming of amino acid biosynthetic gene expression to alleviate nutrient depletion.
基因ID	10985
基因名	GCN1
Swiss	Q92616
别名	GCN1;GCN1L;GCN1L1;PRIC295

## 产品验证



Western blot analysis of GCN1L1 expressed in 293T using GCN1L1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)