

ECH1 Rabbit pAb

货号: **AYP21479**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	35kDa
实测分子量	32KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Mitochondrion,Peroxisome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-328 of human ECH1 (NP_001389.2).
序列	MAAGIVASRRLLDLLTRRLTGSNYPLSISLRLTGSSAQEEASGVALGEAPDHSYSLRVTSAPKQKHLHVQLNRPNKRNAM NKVFWREMVCEFNKISRDCRAVVISGAGKMFTAGIDLMDMASDILQPKGDDVARISWYLRDIITRYQETFNVIERCPKP VIAAVHGGCIGGGVDLVTACDIRYCAQDAFFQVKEVDVGLAADVGTQRLPKVIGNQSLVNELAFTARKMMADEALGSGL VSRVFPDKEVMLDAALALAAEISSKSPVAVQSTKVNLLYSRDHVAESLNYVASWNMSMLQTQDLVKSVQATTENKELKT VTFSKL

靶点信息

研究背景	This gene encodes a member of the hydratase/isomerase superfamily. The gene product shows high sequence similarity to enoyl-coenzyme A (CoA) hydratases of several species, particularly within a conserved domain characteristic of these proteins. The encoded protein, which contains a C-terminal peroxisomal targeting sequence, localizes to the peroxisome. The rat ortholog, which localizes to the matrix of both the peroxisome and mitochondria, can isomerize 3-trans,5-cis-dienoyl-CoA to 2-trans,4-trans-dienoyl-CoA, indicating that it is a delta3,5-delta2,4-dienoyl-CoA isomerase. This enzyme functions in the auxiliary step of the fatty acid beta-oxidation pathway. Expression of the rat gene is induced by peroxisome proliferators.
基因ID	1891
基因名	ECH1
Swiss	Q13011
别名	ECH1;HPXEL

产品验证



Western blot analysis of ECH1 expressed in 293T using ECH1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn