

# PACSIN2 Rabbit pAb

货号: **AYP21543**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	51kDa/55kDa
实测分子量	62KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cell membrane,Cell projection,Cytoplasm,Cytoplasmic side,Cytoplasmic vesicle membrane,Early endosome,Membrane,Peripheral membrane protein,Recycling endosome membrane,caveola,cytoskeleton,ruffle membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 240-410 of human PACSIN2 (NP_009160.2).
序列	RLRFFREVLLLEVQKHLDSLNVAGYKAIYHDLEQSIIRAADAVEDLRWFRANHGPGMAMNWPQFEWSADLNRTLSRREKK KATDGVTLTGINQTDGQSLPSKPSSTLNVPSPNAQSAQSQSSYNPFEDDDTGSTVSEKDDTKAKNVSSYEKTSYPTD WSDDESNNPFSST

## 靶点信息

研究背景	This gene is a member of the protein kinase C and casein kinase substrate in neurons family. The encoded protein is involved in linking the actin cytoskeleton with vesicle formation by regulating tubulin polymerization. Alternative splicing results in multiple transcript variants.
------	---

基因ID	11252
基因名	PACSIN2
Swiss	Q9UNF0
别名	PACSIN2;SDPII

## 产品验证



Western blot analysis of PACSIN2 expressed in HeLa using PACSIN2 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)